



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## PROBLEMS FOR SOLUTION.

### ALGEBRA.

324. Proposed by R. D. CARMICHAEL, Princeton University.

Sum the *finite* series

$$\frac{16n^2 - 2^2}{4!} - \frac{(16n^2 - 2^2)(16n^2 - 4^2)}{6!} + \frac{(16n^2 - 2^2)(16n^2 - 4^2)(16n^2 - 6^2)}{8!} - \dots$$

where  $n$  is a positive integer.

325. Proposed by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

I have a chronometer whose rate is uniform. When it indicates  $t_1$  time at Washington I find that it is  $h_1$  hours slow. I take it to Philadelphia and when it indicates  $t_2$  time, the local time of Philadelphia is  $h_2$  hours faster. I bring my chronometer back to Washington and find that when it indicates  $t_3$  time it is  $h_3$  hours slow. If  $t_1 = 5$  A. M.,  $t_2 = 7$  hours, 54 minutes,  $t_3 = 11$  hours, 46 minutes A. M.,  $h_1 = 1$  hour,  $h_2 = 1 \frac{203}{900}$  hours,  $h_3 = 1 \frac{7}{30}$  hours, find the difference of longitude between Washington and Philadelphia.

### GEOMETRY.

351. Proposed by L. E. DICKSON, Ph. D., The University of Chicago.

Given an isosceles right triangle with hypotenuse  $h$ ; an isosceles triangle with two sides  $h$  and two angles  $A = 22^\circ 30'$ ; a right angle triangle with the same angle  $A$  and opposite side  $h/\sqrt{2}$ ; a triangle with the same angle  $A$ , opposite side  $h$ , and an angle  $45^\circ$ . Form a triangle whose four pieces are these four triangles, and prove geometrically that it is isosceles.

352. Proposed by G. I. HOPKINS, Professor of Astronomy, High School, Manchester, N. H.

Required, to construct the triangle, having given the base, vertical angle and sum of the altitude and the two remaining sides.

353. Proposed by L. H. McDONALD, M. A., Ph. D., Sometimes Tutor at Cambridge, Jersey City, N. J.

In a given circle place two chords which shall be in a given ratio and also a given distance apart.

### CALCULUS.

282. Proposed by S. G. BARTON, Ph. D., Clarkson School of Technology, Potsdam, N. Y.

A rectangular beam of length  $l$  and width  $w$  is taken horizontally from a hall of width  $b$  into a corridor at right angles to the hall. Find the width of the smallest corridor into which it can be taken.

283. Proposed by B. F. FINKEL, Ph. D., Drury College.

By means of the calculus, determine the angle of minimum deviation of a ray of monochromatic light in passing through a triangular prism.